

In this issue of the *Annals of Family Medicine*

Frequency and Prioritization of Patient Health Risks 505

A structured assessment identifies many health risks, but most patients are not ready to change behaviors or discuss health risks with their clinician.

Proactive Case-finding for Older People 514

Geriatric functional screening in New Zealand increases residential placement but does not affect hospitalization or subsequent disability.

Behavioral and Mental Health Assessment in Primary Care 525

Implementing a structured health risk appraisal requires local practice adaptations and about 30 minutes of staff time per patient.

Impact of Continuity on Mortality and Costs 534

In a large nationally representative study of patients in Korea with newly diagnosed hypertension, lower continuity of care is associated with worse cardiovascular outcomes and increased mortality.

Medical Student Debt and Primary Care Practice 542

Student debt has a complex relationship to likelihood of entering primary care practice, varying by type of school.

Techniques for Easier Pill Swallowing 550

Two techniques improve the ease of swallowing tablets and capsules for those with and without swallowing difficulties.

Patient-controlled Taping for Ingrown Toenails 553

Patient-applied taping is a safe and effective nonsurgical method for treatment of ingrown toenails.

Breast Cancer Screening During Hospital Stays 556

Hospitalized women who are overdue for mammograms value the opportunity to have breast cancer screening during their hospital stay.

Validation of Continuity-of-care Measures 559

A measure of year-to-year continuity of care from multiple clinicians captures patients' perceived coordination of care, but is not strongly associated with relational continuity.

Health Information Technology to Keep Patients Insured 568

Health information technology offers opportunities to support health insurance enrollment and retention efforts in primary care practices.

From Triple to Quadruple Aim 573

The Triple Aim should be expanded to include the goal of improving the work life of clinicians and staff.