

In this issue of the *Annals of Family Medicine*

Rates of Uninsured Safety Net Clinic Visits 10

Safety net clinics in states that expanded Medicaid in 2014 had a substantial drop in uninsured visits and an increase in Medicaid-covered visits.

Health Information Technology in Low-Resource Practices 17

Continuing on the “meaningful use” pathway may prove very difficult for many small or low-resource practices.

Office-Based Treatment of Opioid Use Disorder 23

Physicians waived to prescribe buprenorphine-naloxone are largely concentrated in urban areas, which limits access to effective treatment in rural areas.

General Practitioners Recognizing Alcohol Dependence 28

General practitioners identify about the same number of people with alcohol dependence as a structured diagnostic interview, but identify different people.

The Year Before Hospice 33

Functional decline prior to hospice care follows distinct trajectories that can inform optimal palliative care use.

Fixed Ratio vs Lower Limit of Normal of FEV_1/FVC in COPD 41

In diagnosing COPD, combined spirometric criteria of low FEV_1 plus a low ratio of FEV_1/FVC is superior to either criterion alone for predicting clinical outcomes.

Laryngeal Measurements and Diagnostic Tools for COPD 49

Combining lung function questionnaire and laryngeal height can help confirm or discount chronic obstructive pulmonary disease.

Targeting Shoulder Pain 53

Ultrasound imaging enables family physicians to target treatment in patients with acute shoulder pain.

Psychological Treatments for Depression in Primary Care 56

Psychological treatments for adult depression in primary care are effective compared to usual care or placebo, according to this meta-analysis of RCTs.

Pharmacological Treatments for Depression in Primary Care 69

A meta-analysis of RCTs conducted in primary care shows best evidence for use of TCAs and SSRIs in treatment of depression.

Clinical Practice Guideline Executive Summary 80

An AAFP guideline recommends discussing individualized risks and benefits with pregnant women with a history of cesarean births who are deciding between LAC/VBAC and repeat cesarean.

Shine the Light on Direct-to-Consumer Advertising 82

Greater transparency in direct-to-consumer pharmaceutical advertising, including digital forms, could complement data from “Sunshine” regulations.