

In this issue of the *Annals of Family Medicine*

Prediabetes in Healthy-Weight Adults 304

Prediabetes and abdominal obesity have increased among individuals with BMI in the "healthy" range.

Lifestyle Intervention for Obese, Sedentary Patients 311

A home-based intervention for obese patients promotes weight loss and physical activity, but effects diminish after a year.

Association Between Hydration, BMI, and Obesity 320

In a nationally representative sample, inadequate hydration is associated with obesity and elevated BMI.

Diagnosis and Treatment of UTI in Young Children 325

A clinical decision rule based on signs and symptoms can help clinicians identify children who require urinalysis and antibiotic therapy.

Coherence and Adverse Health Outcomes 337

A greater sense of coherence is associated with better health outcomes in people with multimorbidity.

When Do Primary Care Physicians Retire? 344

A study of AMA Masterfile data shows primary care physicians tend to retire in their mid 60s.

Interconception Care by Family Physicians at Well Child Visits 350

Family physicians provide interconception care at well-child visits, and mothers are receptive to the advice.

Seasonality of Ankle Swelling 356

Google Trends data demonstrate seasonality of ankle swelling.

Best-Worst Scaling for Patient Priorities 359

A "best-worst" survey method reveals homeless women's preferences about Papanicolaou test services.

Screening Men for Osteoporosis 365

The Male Osteoporosis Risk Estimation Score is a simple and sensitive way to screen for risk of osteoporosis in men.

Concept Mapping to Engage Patients in CQI 370

Concept mapping can be a method for engaging patients in primary care practice improvement.

Role of Clerks in the PCMH 377

The unique role of clerks can be leveraged to better deliver patient-centered primary care.

A Dance With Mrs Chan 380

A medical student learns that a patient is more than a disease.