

In this issue of the *Annals of Family Medicine*

Clinician Friendly Tool to Identify Clinical Practice Guidelines 413

A simple checklist can help clinicians identify useful clinical practice guidelines.

PCP Workload Based on EHR Log Data and Observation 419

Primary care physicians spend more than half of their work-day interacting with electronic health records during and after clinic hours.

Impact of Scribes on Physician/Patient Satisfaction and Charting Efficiency 427

Scribes improve overall physician satisfaction and charting efficiency, without detracting from patient satisfaction.

Uninsured Primary Care Visits Under the Affordable Care Act 434

Under the Affordable Care Act, Medicaid expansion contributed to sizeable decreases in rates of uninsured visits to community health centers, but disparities remain.

Support for Smoking Cessation After Cancer 443

In UK primary care practices, cancer patients receive less support to quit smoking than patients with heart disease.

Risk Stratification and Care Management in CPC Initiative Practices 451

Practices in a primary care initiative identified more high risk patients and linked more of them to case management using clinician intuition than other methods.

Nature of Blame in Primary Care Patient Safety Incidents 455

In describing primary care patient safety incidents, reporters often attribute blame to the actions of other staff members.

Emergency Hospitalization in Adults With Intellectual Disability 462

Adults with intellectual disabilities are at high risk of preventable emergency hospital admissions.

Developmental Disabilities and Health Care Disparities 471

Individuals with developmental disabilities experience disparities in health status, quality, utilization, access, and unmet health care needs.

Agriculture Health Sectors Collaborate for Population Health 475

Agriculture and health sectors can work together to address communities' health needs.