

Annals of Family Medicine, Inc. 11400 Tomahawk Creek Parkway Leawood, KS 66211-2680 NONPROFIT ORG US POSTAGE PAID LAWRENCE, KS PERMIT NO. 116

#### In this issue of the Annals of Family Medicine

#### Top 20 POEMs of the Past 20 Years 436

Over time, POEMs illustrate the breadth of practice change in primary care and the need for a systematic means of keeping up with the medical literature.

# Professional Loneliness and the Loss of the Doctors' Dining Room 461

Structural changes in medicine, including the demise of the doctors' hospital dining room, have led to an increased sense of professional loneliness.

## Social Isolation and Patient Experience in Older Adults 393

Social isolation is associated with a negative patient experience in older primary care patients in Japan.

#### Adoption of EHR Tools for Social Determinants of Health 399

Systematic EHR-based documentation of social determinant. of health may be feasible, but substantial barriers to adoption exist.

# Opioid-Prescribing Continuity and Risky Opioid Prescriptions 440

Higher prescribing continuity is associated with lower rates of risky opioid prescribing and adverse patient outcomes.

## ED Use and After-Hours Care in a Medical Home 419

Enrollment in a medical home required to provide after hours care is associated with higher emergency department utilization.

# Employment Interventions in Health Settings: Systematic Review and Synthesis 447

To help patients find employment, health care organizations engage employers and use tailored multidisciplinary interpentions

### Predicting an Unfavorable Course of Dizziness in Older Patients 428

A risk score identifies patients at high risk of an unfavorable course of dizziness.

#### Buprenorphine Provision by Early Career Family Physicians 443

Few early career family physicians report being adequately trained in residency to provide buprenorphine treatment, and even fewer provide it.

#### Excess Free Fructose Beverages and Allergy in Children and Adolescents 408

This study provides preliminary evidence for the hypothesis that children with heavy consumption of beverages with excess fructose are more likely to develop allergic symptoms.