

In this issue of the *Annals of Family Medicine*

Older Patients With Dizziness 100

This 10-year study of older primary care patients found that the prognosis and survival rate for patients with dizziness primarily caused by peripheral vestibular disease was better than for those with cardiovascular disease.

Engaging Patients in Advance Care Planning in Outpatient Settings 110

Patient engagement in advance care planning increased after use of an online program at home.

Predicting Opioid Use Following Discharge After Cesarean Delivery 118

24 hours prior-to-discharge opioid intake can help inform opioid prescribing for women after Cesarean delivery.

General Practitioners Compared With Family Physicians in US Medical Practice 127

General practitioners in the United States are a varied group that differs from ABFM-certified family physicians, and the 2 groups should not be combined in research studies.

Anticoagulants' Safety and Effectiveness in General Practice 131

This observational study of oral anticoagulants in French primary care practices observed a lower incidence of overall bleeding with Vitamin K antagonists (VKA) after 1 year, but a 2-times higher incidence of death in the VKA group compared to direct oral anticoagulants (DOACs).

Efficacy and Safety of a Fasting Algorithm 139

Fasting during Ramadan is safe for Muslims with type 2 diabetes who collaborate with their family doctor on managing their diabetes medication.

Polypharmacy Involving Anticholinergics 148

Anticholinergic burden is associated with adverse outcomes in a middle-older aged population as well as in older people.

Maternity Care and Buprenorphine Prescribing in New Family Physicians 156

This analysis, completed using ABFM graduate surveys, characterizes the early-career family physicians who provide maternity care and also prescribe buprenorphine and shows that they are trained in a small number of residency programs.

Impacts of Operational Failures 159

Critical interpretive review of operational failures in primary care finds suboptimal work systems likely have important implications for functional primary care systems.

Voices of Patients With Obesity 169

Patients with obesity share the dread they feel when offering insights to their doctor into the type of help they need.

Co-Design Pacific Style 172

A community advisory group and practice-based research network using the principles of co-design and the Samoan research framework fa'afaletui, generate and answer research questions aimed at decreasing health disparities of Pacific people in South Auckland.

Ethics of Humanitarian Work 176

The ethics of providing humanitarian assistance, despite its significant shortcomings in the midst of epidemics and other conditions among Rohingya refugees, is in refusing to abandon others in desperate situations and in attempting to rectify injustices through witnessing and solidarity.