

Applying Lessons From Behavioral Health Integration to Social Care Integration in Primary Care

Theresa J. Hoeft, PhD¹

Danielle Hessler, PhD²

Damon Francis, MD³

Laura M. Gottlieb, MD, MPH²

¹Department of Psychiatry and Behavioral Sciences, University of Washington, Seattle, Washington

²Department of Family and Community Medicine, University of California San Francisco, San Francisco, California

³Health Leads, Oakland, California

ABSTRACT

Interest and incentives are increasing around strategies whereby the health care sector can better identify and address patients' social and economic needs in the context of primary care delivery. This interest is likely to accelerate during the economic recession following the COVID-19 pandemic. Yet effective and sustainable strategies for integrating social care practices (eg, patient-facing social risk screening and activities to address identified needs) have not been clearly established. Lessons learned from more than 2 decades of research on behavioral health integration could be applied to efforts to integrate social care into primary care. In this article, we synthesize learnings from primary care and behavioral health care integration, and translate them into organizing principles with the goal of advancing social care integration practices to improve the health of both patients and communities.

Ann Fam Med 2021;19:356-361. <https://doi.org/10.1370/afm.2688>.

INTRODUCTION

Growing recognition that patients' social and economic needs influence health outcomes has led to new health care sector initiatives to identify and address those needs in the context of primary care delivery. This work is occurring both in the United States and globally as the importance of integrating health care and social care gains considerable attention.¹ The combination of the COVID-19 pandemic and its economic sequelae has accelerated interest in more systematically integrated care,²⁻⁴ including expanding efforts to screen for social risks⁵⁻⁷ and to leverage health care encounters to intervene on these risks.⁸⁻¹⁰ In some settings, health care systems are using technology to increase capacity to provide social care,¹¹ while in others, systems are also developing and expanding staff capacity (eg, using community health worker models).¹²⁻¹⁵ Emerging research suggests that patient navigation and referral programs can improve outcomes of health and health care use in selected settings.^{8,9,16}

As efforts expand to better integrate social and medical care delivery, it is not surprising that implementation barriers have emerged. A recent National Academies of Sciences, Engineering, and Medicine report has highlighted the growing focus nationally on social care integration alongside looming implementation barriers, including lack of an adequately trained interdisciplinary workforce; poor infrastructure for inter-agency communication and data exchange; and financing models that fail to incentivize interdisciplinary care.¹⁷ These challenges may sound familiar to those who have tracked the progress of behavioral health integration (BHI) in primary care over the last 2 decades. In fact, similar challenges have been faced, and in some cases, gradually dismantled, by BHI advocates across the United States, thereby providing opportunities to learn from this integration in terms of both successes and ongoing limitations. In this article, we identify 3 key lessons learned from BHI successes and challenges that are likely to help advance social care integration activities.

Conflicts of interest: authors report none.

CORRESPONDING AUTHOR

Theresa J. Hoeft
Box 356560
Seattle, WA 98195-6560
thoeft@uw.edu

LESSONS LEARNED FROM BHI

BHI is a systematic collaborative approach to providing patient-centered care using a team of behavioral health and primary care clinicians.¹⁸ Since the Surgeon General's pivotal report in 2001 on integration of mental health into primary care,¹⁹ BHI practices have expanded to include a wide range of behavioral health programs provided in primary care practices; in some places, BHI now also involves providing primary care in behavioral health contexts. Overcoming challenges to BHI has required institutionalizing specific implementation building blocks. Three building blocks have played an outsized role in advancing BHI and are poised to inform social care: (1) consensus on behavioral health metrics and standards that can be used to support incentives for mental health screening and treatment; (2) team-based care facilitators (including colocated services and technology to support shared care for patients and team communication); and (3) context-appropriate adaptation. We discuss each of these building blocks in greater detail below and propose ways in which they could be translated to social care.

Establish Common Metrics, Related Medical Codes, and Incentives for Behavioral Health Needs Assessments and Treatment

BHI Context

The development of the 9-item Patient Health Questionnaire (PHQ-9) brief depression symptom measure in the late 1990s streamlined depression symptom measurement, enabling ready uptake of screening in fast-paced primary care settings.^{20,21} The 2-item Patient Health Questionnaire (PHQ-2) further accelerated depression screening. The importance of a unified, codifiable measure in the field of depression, specifically, cannot be overstated. The PHQ-9 became widely used to compare clinic quality, initially only for screening but subsequently also for tracking and monitoring clinical care.²² Reliance on these measures helped to garner pay-for-performance dollars that could facilitate other BHI infrastructure.

The implementation of billing codes and quality measures in BHI was gradual, but over time, these codes greatly enabled depression care and panel management activities. Early quality metrics for mental health screening were not as popular as physical health metrics (eg, metrics for diabetes) perhaps because mental health metrics were considered more intractable.²³ To incentivize mental health management activities including specialty referrals, however, during 2016–2018 the National Committee for Quality Assurance added Healthcare Effectiveness Data and Information Set quality metrics for having (1) PHQ-9 score documented during follow-up visits for patients with major

depressive disorder or dysthymia; (2) documented remission or response within 4 to 8 months of an initially elevated PHQ-9 score; and (3) follow-up visits scheduled within 30 days if PHQ-9 screening was initially positive.²²

Use of common metrics in BHI is not limited to depression. Other common measures have facilitated uptake and quality measure development for Screening, Brief Intervention, and Referral to Treatment programs, which focus on assessment for and treatment of substance use disorders.^{24,25} As an example, the National Committee for Quality Assurance added a Healthcare Effectiveness Data and Information Set quality measure to track alcohol use and referrals; the committee also developed a learning collaborative to support health plans in adopting this measure.²⁶ The availability of Current Procedural Terminology billing codes also helped to facilitate reimbursement for Screening, Brief Intervention and Referral to Treatment screening and other behavioral health interventions in primary care.²⁷

Translation to Social Care

There is little consensus on effective, valid social risk–screening tools in the United States. A stronger commitment to developing a common core of key social risk measures may speed the development of screening adoption and facilitate development of payment or reimbursement incentives as it did for depression and substance use screening. Medical codes that bridge measures across screening tools—for example, measures of food insecurity from different screening tools—could be used as an alternative while consensus on a common core of measures is developed.²⁸ Implementation research can help to establish how to use measures (eg, focusing on high-risk target populations, using electronic health record modules).

Common metrics can enable standardization of social care interventions and simultaneously provide data needed to gauge population-level initiatives related to social conditions. In depression treatment, standardization of measures was effective in part because the related incentives were not isolated to screening—they also incorporated strategies to promote panel management and follow-up mental health care, as well as provision of limited social care in cases where expertise aligns. Medical codes and payment models in the social care space will need to similarly ensure that common metrics not only increase screening practices, but also contribute to improved social health outcomes for individuals and populations. Possible sources of funding to support this social care work include Advanced Payment Models and Medicare Advantage plans.^{29–31} Most recently, the American Medical Association and the Centers for Medicare and

Medicaid Services have set forth changes in the 2021 Evaluation and Management medical coding guidelines to ensure clinicians can account for their work with socially complex patients in fee-for-service billing.³²

Reinforce Comprehensive Team-Based Primary Care and Strong Community Linkages

BHI Context

An early lesson in BHI was that even with incentives, clinicians often believed that screening in the absence of effective treatment teams was untenable.^{33,34} As a result, efforts to reinforce comprehensive, team-based primary care and strong community linkages to mental health supports were critical facilitators for successful BHI. Primary care integration helped to decrease delays in mental health consultations and increased treatment adherence because patients were able to be seen in settings that were more accessible and potentially less stigmatizing. Colocation provided new opportunities to share practice culture across clinician groups.¹⁸

Improving BHI also required other team-based care reinforcements, including those related to technology. Technology was used to strengthen communication and cohesion among the team (eg, use of cell phones and instant messaging to communicate; data sharing and tracking of patient populations). Important examples of technology support in BHI included use of patient registries to support team-based patient reviews; ability of all team members to perform patient scheduling; and access to mental and physical health electronic health records for the entire team. Registries, particularly, continue to help teams organize and monitor consultation, education, and support services; follow changes in treatment; and track symptoms and outcomes,³⁵ especially in cases where primary care and behavioral health electronic health records are not seamless.

Translation to Social Care

As in behavioral health care, some socioeconomic barriers to health can be overcome through low-touch social care support from the primary care team, but many patients facing social and economic challenges will require more intensive and/or longer-term assistance from organizations equipped to provide social services, and just as importantly, collective advocacy to improve community social conditions. Challenges include determining which patients will benefit from differing levels and types of care, designing complementary programs that can support patients with specific needs, and developing robust partnerships and advocacy agendas to ensure needs are met both initially and over time. As in BHI, achieving these goals

may require colocation of services, stepped models of care, workforce development (eg, training everyone on the team including front desk staff on addressing social needs), and increased reliance on technology to facilitate partnerships. The team should include someone with expertise in social care (eg, social workers).¹⁷ As in BHI, this integration work will demand working through cultural differences across organizations³⁶ and possibly between the social care expert and the primary care team. Acknowledging these differences and working toward organizational integration of cultures can shift practice to advance whole-person care.

An added barrier to organizational alignment in social care integration is that there are few payment models that bridge clinical and nonclinical systems providing care. These models will need to be developed and scaled to facilitate effective integration. As with BHI, however, these efforts will be worthwhile because addressing social care needs in primary care and in partnership with community partners can improve access to services and destigmatize their uptake. In addition, by asking about social needs, clinicians may both build rapport with patients and help tailor their care to those needs (eg, offering nonrefrigerated medications to patients who are homeless).

Technology also will be needed to support cross-disciplinary, team-based care. This technology might involve direct communication modalities (eg, use of cell phones and instant messaging) and registries, but it also will need to expand to interagency data-sharing infrastructure. Innovative examples include San Diego County's 211 Community Information Exchange, which integrates key client data across medical and social institutions³⁷ and other community referral resource platforms, with the aim of increasing inter-organizational communication and improving patient care and population health planning.¹¹

Enable Evidence-Based Evolution and Context-Appropriate Adaptation

BHI Context

A third important aspect of BHI success involved using context-dependent adaptation to increase model adoption and improve care for specific populations. Although standardization of behavioral health measures was helpful to facilitate integration, implementing and scaling programs in clinic settings has also required that core models maintain flexibility to clinic context. One of the core models of primary care-based BHI that has been successfully adapted for use in different settings, for instance, is the collaborative care model (CoCM) based on the chronic care model.³⁸ The CoCM foundation includes a care manager, who offers both clinic visits and telephone

outreach; a psychiatric consultant; and a registry to track measurement-based care for all patients with diagnosable depression or anxiety.^{35,39,40} This model has been successfully adapted to a range of health care delivery contexts.⁴¹⁻⁴³ The 5 principles of CoCM⁴⁴ are present in each of these adaptations, which differ in terms of staffing, setting, and condition of focus. A second core model for BHI is the behavioral health consultant model, which integrates these consultants into primary care clinics.⁴⁵ Some settings have successfully merged the CoCM and behavioral health consultant models to improve care for patients with varying levels of behavioral health needs.⁴⁶ Adaptations have involved preimplementation needs assessments that helped clinics overcome context-specific barriers and identify key unique facilitators of integrated activities.^{47,48} The Exploration, Preparation, Implementation, Sustainment framework in implementation science has been applied in efforts to adapt the CoCM to different BHI contexts.^{49,50}

Translation to Social Care

Social care interventions are frequently dependent on the community social resources landscape. Implementation will therefore depend on each local context, including unique population needs, community resources, partners, and other stakeholders.⁵¹⁻⁵⁴ Partnerships may include connections with local public health agencies and other organizations that can attend to both structural and policy drivers that are shaping patients' social conditions. Such partnerships can strengthen the availability of social care resources for both individual patients and communities. Involving these partners in the preimplementation needs assessment stage would thus improve both implementation and sustainability of social care interventions. Collectively, health care systems, public health agencies, and local community-based organizations can advocate for needed services in their area. Advocacy efforts such as this could in turn offer insights back to BHI. For instance, BHI advocacy is needed to address mental health resource shortages in rural and low-resource settings and to improve outreach and care for opioid use disorders or suicide risk.

BHI has also typically focused on individuals rather than connections to broader coalitions working on public health. Both BHI and social care integration can learn from larger campaigns such as antismoking efforts that resulted in actions by a broad coalition including health care. Universal screening and education allowed those in health care to support efforts with individuals *within* a broader strategy for policy and institutional changes that can have a much greater public health impact, as we have seen with antismoking efforts.

CONCLUSIONS

In 2008, the World Health Organization issued a report outlining a 7-point rationale for the integration of mental health and primary care services.⁵⁵ Many of the arguments they articulated translate easily into a rationale for better integrating social care into primary care. For example, the burden of social disadvantage is large; social and physical health problems are interwoven; an important and substantial treatment gap exists; primary care can increase access to treatment; and using primary care for social care can promote respect for basic human rights. There is also emerging evidence to suggest that social care integration activities can improve health⁵⁶⁻⁵⁹ and decrease health care costs.⁶⁰⁻⁶⁴

As programs on social care integration develop, we can and should learn from the BHI movement. Applying BHI's hard-won lessons would mean developing core metrics and incentives, directing supports to the diverse team needed to facilitate interventions related to social and economic needs, and improving implementation and scaling efforts by enabling contextualization. Just as importantly, we should learn from the limited impact BHI has had on behavioral health outcomes at the population level, manifested most strikingly in the ongoing epidemics of suicides⁶⁵ and drug overdoses.⁶⁶ The effectiveness of primary care programs targeting both behavioral and social factors to improve health outcomes is ultimately more dependent on local, state, and federal policies that shape individual and community health than on health care-specific activities. Indeed, as BHI and social care integration efforts advance, their greatest impact may lie in deepening our collective commitment to ensuring that all people have the resources and relationships they need to attain and sustain good health.

To read or post commentaries in response to this article, go to <https://www.AnnFamMed.org/content/19/4/356/tab-e-letters>.

Key words: behavioral health integration; social care integration; social and economic needs; social determinants of health; social context; health services; population health; vulnerable populations

Submitted July 29, 2020; submitted, revised, October 26, 2020; accepted December 26, 2020.

Funding support: Dr Gottlieb's work on this article was supported by grants from the Robert Wood Johnson Foundation and Kaiser Permanente.

Disclaimer: The views expressed are solely those of the authors and do not necessarily represent official views of the authors' affiliated institutions or funders.

References

1. Bhattacharyya O, Shaw J, Sinha S, et al. Innovative integrated health and social care programs in eleven high-income countries. *Health Aff (Millwood)*. 2020;39(4):689-696.

2. Brewster AL, Fleming M, Knox M, Roth A. Community resilience for COVID-19 and beyond: leveraging a system for health care and social services integration. Published Aug 12, 2020. Accessed Oct 24, 2020. <https://www.healthaffairs.org/doi/10.1377/hblog20200807.222833/full>
3. Fichtenberg C, Gottlieb LM. Health and social services integration is mission-critical in the coronavirus response. Published Apr 12, 2020. Accessed Oct 24, 2020. <https://www.healthaffairs.org/doi/10.1377/hblog20200414.50259/full>
4. Charlton G. Integration and collaboration during COVID-19. Health and Social Care Alliance Scotland. Published Jan 5, 2020. Accessed Oct 24, 2020. <https://www.alliance-scotland.org.uk/blog/opinion/integration-and-collaboration-during-covid-19/#expanded>
5. Garg A, Cull W, Olson L, et al. Screening and referral for low-income families' social determinants of health by US pediatricians. *Acad Pediatr*. 2019;19(8):875-883.
6. Frazee TK, Brewster AL, Lewis VA, Beidler LB, Murray GF, Colla CH. Prevalence of screening for food insecurity, housing instability, utility needs, transportation needs, and interpersonal violence by US physician practices and hospitals. *JAMA Netw Open*. 2019;2(9):e1911514.
7. Andermann A. Screening for social determinants of health in clinical care: moving from the margins to the mainstream. *Public Health Rev*. 2018;39:19.
8. Gottlieb LM, Wing H, Adler NE. A systematic review of interventions on patients' social and economic needs. *Am J Prev Med*. 2017;53(5):719-729.
9. Fichtenberg CM, Alley DE, Mistry KB. Improving social needs intervention research: key questions for advancing the field. *Am J Prev Med*. 2019;57(6 Suppl 1):S47-S54.
10. Alderwick HAJ, Gottlieb LM, Fichtenberg CM, Adler NE. Social prescribing in the U.S. and England: emerging interventions to address patients' social needs. *Am J Prev Med*. 2018;54(5):715-718.
11. Cartier Y, Fichtenberg C, Gottlieb L. *Community Resource Referral Platforms: A Guide for Health Care Organizations*. Published 2019. <https://sirenetwork.ucsf.edu/tools-resources/resources/community-resource-referral-platforms-guide-health-care-organizations>
12. Hsu C, Hertel E, Johnson E, et al. Evaluation of the Learning to Integrate Neighborhoods and Clinical Care project: findings from implementing a new lay role into primary care teams to address social determinants of health. *Perm J*. 2018;22.
13. Zeigler BP, Redding SA, Leath BA, Carter EL. Pathways community HUB: a model for coordination of community health care. *Popul Health Manag*. 2014;17(4):199-201.
14. Redding S, Conrey E, Porter K, Paulson J, Hughes K, Redding M. Pathways community care coordination in low birth weight prevention. *Matern Child Health J*. 2015;19:643-650.
15. Kangovi S, Mitra N, Norton L, et al. Effect of community health worker support on clinical outcomes of low-income patients across primary care facilities: a randomized clinical trial. *JAMA Intern Med*. 2018;178(12):1635-1643.
16. Kangovi S, Mitra N, Grande D, Long JA, Asch DA. Evidence-based community health worker program addresses unmet social needs and generates positive return on investment. *Health Aff (Millwood)*. 2020;39(2):207-213.
17. National Academies of Sciences, Engineering, and Medicine. *Integrating Social Needs Care Into the Delivery of Health Care to Improve the Nation's Health*. Published 2019. Accessed Oct 24, 2020. <https://doi.org/10.17226/25467>
18. Peek C, National Integration Academy Council. *Lexicon for Behavioral Health and Primary Care Integration: Concepts and Definitions Developed by Expert Consensus*. Published 2013. Accessed May 20, 2021. <https://integrationacademy.ahrq.gov/sites/default/files/2020-06/Lexicon.pdf>
19. Department of Health and Human Services (DHHS). *Report of a Surgeon General's Working Meeting on the Integration of Mental Health Services and Primary Health Care*. Published 2001. Accessed Oct 24, 2020. <https://www.ncbi.nlm.nih.gov/books/NBK44335/>
20. Spitzer RL, Kroenke K, Williams JB. Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. *Primary Care Evaluation of Mental Disorders. Patient Health Questionnaire. JAMA*. 10 1999;282(18):1737-1744
21. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med*. 2001;16(9):606-613.
22. National Committee for Quality Assurance (NCQA). HEDIS depression measures specified for electronic clinical data systems. Accessed Jun 27, 2020. <https://www.ncqa.org/hedis/the-future-of-hedis/hedis-depression-measures-specified-for-electronic-clinical-data/>
23. Anderson B. HEDIS antidepressant medication management measures and performance-based measures: an opportunity for improvement in depression care. *Am J Manag Care*. 2007;13(4 Suppl):S98-S102.
24. Babor TF, McRee BG, Kassebaum PA, Grimaldi PL, Ahmed K, Bray J. Screening, Brief Intervention, and Referral to Treatment (SBIRT): toward a public health approach to the management of substance abuse. *Subst Abus*. 2007;28(3):7-30.
25. Hargraves D, White C, Frederick R, et al. Implementing SBIRT (Screening, Brief Intervention and Referral to Treatment) in primary care: lessons learned from a multi-practice evaluation portfolio. *Public Health Rev*. 2017;38:31.
26. National Committee for Quality Assurance (NCQA). New HEDIS measure strengthens alcohol screening and follow up. Accessed Jun 27, 2020. <https://www.ncqa.org/news/new-hedis-measure-strengthens-alcohol-screening-and-follow-up/>
27. Substance Abuse and Mental Health Services Administration (SAMHSA). Screening, brief intervention, and referral to treatment (SBIRT). Accessed Jun 27, 2020. <https://www.samhsa.gov/sbirt>
28. SIREN UCSF, Robert Wood Johnson Foundation, EMI Advisors. *The Gravity Project: A Social Determinants of Health Coding Collaborative*. Published 2019. Accessed Oct 24, 2020. <https://confluence.hl7.org/download/attachments/51227176/Gravity%20Use%20Case%20Package%20FINAL%2020190930.pdf>
29. American Academy of Family Physicians. Advancing health equity by addressing the social determinants of health in family medicine (position paper). Published Apr 2019. Accessed Oct 24, 2020. <https://www.aafp.org/about/policies/all/social-determinants-health-family-medicine.html>
30. DuGoff EH, Buckingham W, Kind AJH, Chao S, Anderson GF. *Targeting High-Need Beneficiaries in Medicare Advantage: Opportunities to Address Medical and Social Needs*. Published 2019. Accessed Oct 24, 2020. https://www.commonwealthfund.org/sites/default/files/2019-02/DuGoff_targeting_high-need_beneficiaries_ib_0.pdf
31. Liss S. Americans want their doctors to ask about social needs. Published Jun 5, 2019. Accessed Oct 24, 2020. <https://www.healthcarediver.com/news/americans-want-their-doctors-to-ask-about-social-needs/556197/>
32. Deighan C. Social determinants of health and 2021 E&M code changes. Published Mar 30, 2020. Accessed Oct 24, 2020. <https://www.icd10monitor.com/social-determinants-of-health-and-2021-e-m-code-changes>
33. O'Connor EA, Whitlock EP, Beil TL, Gaynes BN. Screening for depression in adult patients in primary care settings: a systematic evidence review. *Ann Intern Med*. 2009;151(11):793-803.
34. Gjerdingen DK, Yawn BP. Postpartum depression screening: importance, methods, barriers, and recommendations for practice. *J Am Board Fam Med*. 2007;20(3):280-288.
35. Unutzer J, Katon W, Callahan CM, et al. Collaborative care management of late-life depression in the primary care setting: a randomized controlled trial. *JAMA*. 2002;288(22):2836-2845.

36. Catalyz. *Building to a sum greater than its parts: a hands-on guide to cultural integration in community health partnerships and alliances*. Published Oct 15, 2019. Accessed Oct 24, 2020. <https://www.chcf.org/publication/hands-on-guide-cultural-integration-partnerships/>
37. San Diego 211. Accessed Jun 27, 2020. <https://211sandiego.org/>
38. Wagner EH. Chronic disease management: what will it take to improve care for chronic illness? *Eff Clin Pract*. 1998;1(1):2-4.
39. Katon W, Von Korff M, Lin E, et al. Collaborative management to achieve treatment guidelines. Impact on depression in primary care. *JAMA*. 1995;273(13):1026-1031.
40. Katon W, Robinson P, Von Korff M, et al. A multifaceted intervention to improve treatment of depression in primary care. *Arch Gen Psychiatry*. 1996;53(10):924-932.
41. Lyon AR, Whitaker K, French WP, Richardson LP, Wasse JK, McCauley E. Collaborative care in schools: enhancing integration and impact in youth mental health. *Adv Sch Ment Health Promot*. 2016; 9(3-4):148-168.
42. Roy-Byrne P, Craske MG, Sullivan G, et al. Delivery of evidence-based treatment for multiple anxiety disorders in primary care: a randomized controlled trial. *JAMA*. 2010;303(19):1921-1928.
43. Zatzick D, Rivara F, Jurkovich G, et al. Enhancing the population impact of collaborative care interventions: mixed method development and implementation of stepped care targeting posttraumatic stress disorder and related comorbidities after acute trauma. *Gen Hosp Psychiatry*. 2011;33(2):123-134.
44. AIMS Center. Principles of collaborative care. Accessed Jun 27, 2020. <https://aims.uw.edu/collaborative-care/principles-collaborative-care>
45. Reiter JT, Dobmeyer AC, Hunter CL. The primary care behavioral health (PCBH) model: an overview and operational definition. *J Clin Psychol Med Settings*. 2018;25(2):109-126.
46. AIMS Center. Stepped model of integrated behavioral health care. Accessed Jun 27, 2020. <https://aims.uw.edu/stepped-model-integrated-behavioral-health-care>
47. Kirchner JE, Cody M, Thrush CR, Sullivan G, Rapp CG. Identifying factors critical to implementation of integrated mental health services in rural VA community-based outpatient clinics. *J Behav Health Serv Res*. 2004;31(1):13-25.
48. Kirchner JE, Ritchie MJ, Pitcock JA, Parker LE, Curran GM, Fortney JC. Outcomes of a partnered facilitation strategy to implement primary care-mental health. *J Gen Intern Med*. 2014;29(Suppl 4): 904-912.
49. Aarons GA, Hurlburt M, Horwitz SM. Advancing a conceptual model of evidence-based practice implementation in public service sectors. *Adm Policy Ment Health*. 2011;38(1):4-23.
50. EPIS framework. Accessed Jun 27, 2020. <https://episframework.com/>
51. Blachman-Demner DR, Wiley TRA, Chambers DA. Fostering integrated approaches to dissemination and implementation and community engaged research. *Transl Behav Med*. 2017;7(3):543-546.
52. Wallerstein N, Duran B. Community-based participatory research contributions to intervention research: the intersection of science and practice to improve health equity. *Am J Public Health*. 2010; 100(Suppl 1):S40-S46.
53. Eder MM, Carter-Edwards L, Hurd TC, Rumala BB, Wallerstein N. A logic model for community engagement within the Clinical and Translational Science Awards consortium: can we measure what we model? *Acad Med*. 2013;88(10):1430-1436.
54. Wells K, Miranda J, Bruce ML, Alegria M, Wallerstein N. Bridging community intervention and mental health services research. *Am J Psychiatry*. 2004;161(6):955-963.
55. World Health Organization. *Integrating Mental Health Into Primary Care: A Global Perspective*. Published 2008. Accessed Oct 24, 2020. https://whqlibdoc.who.int/publications/2008/9789241563680_eng.pdf
56. Gennetian LA, Miller C. Children and welfare reform: a view from an experimental welfare program in Minnesota. *Child Dev*. 2002; 73(2):601-620.
57. Ryan AM, Kutob RM, Suther E, Hansen M, Sandel M. Pilot study of impact of medical-legal partnership services on patients' perceived stress and wellbeing. *J Health Care Poor Underserved*. 2012;23(4): 1536-1546.
58. Gurvey J, Rand K, Daugherty S, Dinger C, Schmeling J, Laverty N. Examining health care costs among MANNA clients and a comparison group. *J Prim Care Community Health*. 2013;4(4):311-317.
59. Milstein A, Gilbertson E. American medical home runs. *Health Aff (Millwood)*. 2009;28(5):1317-1326.
60. Rosenheck R, Kaspro W, Frisman L, Liu-Mares W. Cost-effectiveness of supported housing for homeless persons with mental illness. *Arch Gen Psychiatry*. 2003;60(9):940-951.
61. Buchanan D, Kee R, Sadowski LS, Garcia D. The health impact of supportive housing for HIV-positive homeless patients: a randomized controlled trial. *Am J Public Health*. 2009;99(Suppl 3): S675-S680.
62. Basu A, Kee R, Buchanan D, Sadowski LS. Comparative cost analysis of housing and case management program for chronically ill homeless adults compared to usual care. *Health Serv Res*. 2012;47(1 Pt 2): 523-543.
63. Herman DB, Conover S, Gorroochurn P, Hinterland K, Hoepner L, Susser ES. Randomized trial of critical time intervention to prevent homelessness after hospital discharge. *Psychiatr Serv*. 2011;62(7): 713-719.
64. Tomita A, Herman DB. The impact of critical time intervention in reducing psychiatric rehospitalization after hospital discharge. *Psychiatr Serv*. 2012;63(9):935-937.
65. Hedegaard H, Curtin SC, Warner M. *Suicide Mortality in the United States, 1999-2017*. NCHS Data Brief, no. 330. Published Nov 2018. Accessed Oct 24, 2020. <https://www.cdc.gov/nchs/products/databriefs/db330.htm>
66. Hedegaard H, Minino AM, M W. *Drug Overdose Deaths in the United States, 1999-2018*. NCHS Data Brief, no. 356. Published Jan 2020. Accessed Oct 24, 2020. <https://www.cdc.gov/nchs/products/databriefs/db356.htm>