

# Agile Implementation of a Digital Cognitive Assessment for Dementia in Primary Care

Diana Summanwar, MD<sup>1</sup>

Nicole R. Fowler, PhD, MHSA<sup>2,3</sup>

Dustin B. Hammers, PhD, ABPP-CN<sup>4</sup>

Anthony J. Perkins, MS<sup>4</sup>

Jared R. Brosch, MD<sup>5</sup>

Deanna R. Willis, MD, MBA<sup>1</sup>

<sup>1</sup>Department of Family Medicine, Indiana University School of Medicine, Indianapolis, Indiana

<sup>2</sup>Department of Medicine, Indiana University School of Medicine, Indianapolis, Indiana

<sup>3</sup>Indiana University Center for Aging Research, Indianapolis, Indiana

<sup>4</sup>Department of Statistics and Data Sciences, Indiana University School of Medicine, Indianapolis, Indiana

<sup>5</sup>Department of Neurology, Indiana University School of Medicine, Indianapolis, Indiana

## ABSTRACT

**PURPOSE** This study aimed to assess how agile implementation–driven iterative processes and tailored workflows can facilitate the implementation of a digital cognitive assessment (DCA) tool for patients aged 65 years or older into primary care practices.

**METHODS** We used agile implementation principles to integrate a DCA tool into routine workflows across 7 primary care clinics. The intervention involved a structured selection process for identifying an appropriate DCA tool, stakeholder engagement through iterative sprints (structured, time-bound cycles), and development of tailored workflows to meet clinic-specific needs. A brain health navigator role was established to support patients with positive or borderline screenings, and assist primary care clinicians with follow-up assessment. We used the reach, effectiveness, adoption, implementation, and maintenance (RE-AIM) framework to evaluate the intervention’s performance over a 12-month period.

**RESULTS** The intervention engaged 69 (63.8%) of 108 clinicians across the 7 clinics. DCA screening was completed in 1,808 (10.8%) of 16,708 eligible visits. We selected the Linus Health Core Cognitive Evaluation tool as our DCA tool based on stakeholder evaluations. Screening workflows were tailored to each clinic. The brain health navigator received 447 referrals for further assessment of a positive or borderline screening result. Four clinics fully adopted the intervention, achieving a DCA completion rate of at least 20%, and 5 clinics were still routinely using the DCA tool at 12 months.

**CONCLUSIONS** Agile implementation effectively helped integrate the DCA tool into primary care workflows. Customized workflows, stakeholder engagement, and iterative improvements were crucial for adoption and sustainability. These insights can guide future efforts for early detection and management of cognitive impairment in primary care, ultimately improving patient outcomes and easing the burden on health care professionals.

*Ann Fam Med* 2025;23:199-206. <https://doi.org/10.1370/afm.240294>

## INTRODUCTION

Alzheimer disease and related dementias affect approximately 6.7 million adults in the United States<sup>1</sup> with cases among underserved populations such as non-Hispanic Black and Hispanic older adults (aged ≥65 years) disproportionately undiagnosed and undertreated.<sup>1-3</sup> This gap in care is concerning given the suggested cost-effectiveness of early detection<sup>4-6</sup> and advances in anti-amyloid therapies for Alzheimer disease.<sup>7-11</sup>

Primary care clinicians play a critical role in early detection<sup>12</sup> but face barriers<sup>13-16</sup> including a lack of validated screening tools embedded in electronic health records, inadequate reimbursement for cognitive assessments, time constraints, and insufficient training and confidence in diagnosing cognitive impairments.<sup>12,17-20</sup>

Limited awareness regarding the benefits of early intervention and modest therapeutic efficacy<sup>21</sup> further impact detection efforts; however, collaborative care<sup>12</sup> has improved quality of life for older adults with mild cognitive impairment, even before the advent of newer therapies.<sup>22</sup> Real-world evidence on integrating screening into outpatient clinic settings is limited, highlighting the need for scalable solutions in primary care workflows.

Implementation science offers frameworks to bridge the research-to-practice gap by addressing barriers to translating evidence into routine care.<sup>23</sup> Agile implementation applies principles from behavioral economics, complex adaptive systems theory, and network science to understand and “nudge” the behaviors of individuals and organizations toward successful implementation.<sup>24,25</sup> It emphasizes

Annals Early Access article

Conflicts of interest: authors report none.

### CORRESPONDING AUTHOR

Diana Summanwar  
1040 Wishard Blvd  
Indianapolis, IN 46202  
[dsummanw@iu.edu](mailto:dsummanw@iu.edu)  
[dsummanw@iuhealth.org](mailto:dsummanw@iuhealth.org)

rapid, iterative development and deployment of interventions according to an agreed-on cross-site minimum standard,<sup>26</sup> allowing for the tailoring of interventions to the unique needs of specific clinical settings, which ensures both effectiveness and sustainability.<sup>24,25</sup> Agile implementation identifies implementation champions<sup>27</sup> locally through engagement of early engagers, a type of clinician persona.<sup>28-30</sup> It has been shown to allow generalizability of a common standardized approach in the context of locally customized implementation in health care settings.<sup>24,25,28,31-34</sup> This study evaluated the implementation of a digital cognitive assessment (DCA) tool into primary care using agile implementation. The objectives were to identify a tool acceptable to stakeholders, develop clinic-specific screening workflows, and optimize care pathways for diagnostic assessment of positive screenings.

Patient navigation in primary care is an essential approach for addressing barriers to care and bridging gaps caused by health care fragmentation.<sup>35</sup> Navigation supports those with complex needs by connecting patients with resources and facilitating care coordination. To address clinician-identified barriers such as limited time, resources, knowledge, and confidence in evaluating patients for cognitive impairment,<sup>12,17-20</sup> we also introduced a brain health navigator (BHN) role.

## METHODS

### Study Overview

We conducted a pragmatic clinical quality improvement project<sup>36-38</sup> using agile implementation to integrate DCA into workflows across 7 diverse primary care clinics. Clinics were selected based on panel size of older adults (aged  $\geq 65$  years), demographic diversity, and variety of practice type and location. We used Area Deprivation Index scores,<sup>39,40</sup> ranging from 1 to 100, to measure neighborhood disadvantage.

We developed and implemented the DCA screening intervention and then monitored outcomes over 12 months. Development had 4 key components: DCA tool selection, primary care leadership and clinician engagement, development of a workflow tailored to each clinic's specific needs, and establishment of the BHN role. We identified clinicians exhibiting early engager characteristics<sup>27</sup> and involved them early in the process.

This study was approved by the Indiana University Institutional Review Board as a quality improvement study (study no. 15325) and an evaluation exempt study (study no. 15281), with a waiver granted for informed consent.

### Study Setting

The study was conducted at 7 Indiana University Health primary care clinics, from June 1, 2022, to May 31, 2023, as part of the Davos Alzheimer's Collaborative Healthcare System Preparedness Early Detection Flagship program (hereafter Davos Flagship program). Their characteristics are shown in [Table 1](#). Of the 7 clinics, 4 were suburban (including 1 value-based clinic) and 3 were urban (including 1 value-based clinic and 1 residency clinic). (A value-based clinic is a health care practice that prioritizes patient outcomes over the volume of services provided. These clinics use team-based care and payment models that reward quality and improved health rather than quantity.)

### Eligibility Criteria for Screening

Patients aged 65 years or older presenting for any primary care visit were eligible for DCA screening; however, to establish realistic goals, engage clinicians and staff meaningfully, and promote incremental improvement, clinicians and clinics were allowed to restrict eligibility to patients this age making specific types of visits. Through collaborative meetings at each clinic, we incorporated clinician and staff input to

**Table 1. Baseline Characteristics of Participating Primary Care Clinics**

Clinic	Location <sup>a</sup>	No. of patients aged $\geq 65$ years <sup>b</sup>	No. of clinicians	Patient race and ethnicity, %		Patient insurance, %				Patient ADI national percentile <sup>c</sup>
				Black/African American	Hispanic/Latino	Medicare	Medicaid	Commercial	None	
1	Suburban	1,690	5	7.1	2.8	37.3	15.1	45.9	1.5	59
2	Suburban	1,930	7	2.3	2.4	37.3	15.1	45.9	1.5	59
3	Urban	314	3	39.0	1.0	42.6	9.3	46.5	1.3	79
4	Suburban	1,697	3	8.7	2.5	90.7	0.1	9.1	0.1	24
5	Urban	1,324	71	51.4	7.1	13.0	50.0	13.0	17.0	66
6	Suburban	1,547	10	5.4	1.7	17.2	10.0	71.3	1.3	24
7	Urban	3,000	9	28.6	3.0	31.7	13.8	52.8	1.4	66

ADI = Area Deprivation Index.

<sup>a</sup> Clinics 3 and 4 were value-based clinics; clinic 5 was an academic medical center clinic with a residency program, located in a Health Professional Shortage Area; and clinic 7 was an academic medical center clinic located in a Health Professional Shortage Area.

<sup>b</sup> Number of unique patients in this age group seen per year.

<sup>c</sup> Possible values range from 1 to 100; higher values indicate greater socioeconomic disadvantage.

develop tailored criteria that aligned with the unique operational characteristics of that clinic. Depending on the clinic, eligible visits were certain types of visits—Annual Medicare Wellness Visits, routine nonurgent follow-up visits, or visits occurring during specific clinic sessions and times—or all visits by patients aged 65 years or older. Patients meeting the eligibility criteria for their clinic were flagged in the electronic health record system, prompting the medical assistant to perform the DCA screening during the visit. When an eligible patient was not screened, we classified the screening as missed, declined by clinician, or declined by patient.

### Agile Implementation Process

The agile implementation process involved 8 key steps, detailed below.

#### 1. Identifying Opportunities for Improvement

We formed an interdisciplinary team to participate in the Davos Flagship program consisting of primary care leaders, specialists in Alzheimer disease and related dementias, agile implementation specialists, DCA tool representatives, clinic physicians, clinical staff champions, and research assistants.

#### 2. Selecting a DCA Tool

We designed a sprint (a structured, time-bound cycle), guided by the Delphi method,<sup>41</sup> to select the DCA tool from those offered by 3 commercial DCA vendors approved by the Davos Flagship program.<sup>42</sup> We defined 3 key stakeholder

groups: research (Alzheimer disease and related dementias researchers); clinical (primary care clinicians and administrative leaders); and patient and medical assistant (Patient and Family Advisory Council members and medical assistants). We used discussions with key informants from each group to develop a list of potential aspects of the DCA tool or DCA workflow that would be of particular interest to that group. Before engaging with DCA vendors, stakeholders completed an online questionnaire to rank-order tool selection criteria based on the key informant list. Weighted rankings of the criteria are shown in [Table 2](#).

Each stakeholder group received vendor information in advance and attended a 2-hour session that included 30-minute vendor presentations followed by question-and-answer segments. Stakeholders then assessed each vendor via an online evaluation, and the weighted decision matrix<sup>43</sup> was used to score each tool against the defined criteria.

#### 3. Developing an Evaluation Plan

We set process and outcome measures to evaluate the implementation. Tracking and reporting strategies were developed and implemented to support the implementation measurement plan.

#### 4. Mapping Current Processes

A dedicated sprint was performed to gauge clinician engagement levels. We conducted preintervention surveys and interactive sessions to assess clinicians' perceptions and preparedness, as well as to make inquiries into

unique aspects of their practices influencing the screening process. The concept of a Minimum Viable Solution prototype was introduced to define essential non-negotiable elements in the screening process. Comprehensive feedback from all stakeholders and analysis of performance metrics were emphasized. We customized evidence-based solutions for each clinic, adjusting clinician schedules, designating medical assistant representatives, and identifying physical spaces for related activities.

#### 5. Conducting Implementation Sprints

We conducted sprints to iteratively test the identified intervention specifications. These sprints included clinician protocol training sessions designed to familiarize clinicians with the DCA tool and its integration into routine practice, and team member DCA training focused on equipping medical assistants with the necessary skills to administer the DCA and manage workflows. In addition, workflow development was iteratively tailored to

**Table 2. Criteria for Digital Cognitive Assessment Tool Selection**

Stakeholder group	Criteria assessed by stakeholder group	Relative weight within stakeholder group criteria <sup>a</sup>
Research	Demonstrated accuracy of primary care physician result interpretation	0.219
	Access to aggregate data reporting for all patients to whom the DCA is administered	0.213
	Demonstrated evidence base for methodologic approach behind the DCA	0.201
	Ability to link DCA results to clinical data	0.189
	Promotes or supports research work across Indiana University ADRD research portfolio	0.177
Clinical	Ease of report acquisition (eg, portal delivery)	0.235
	Ease of setup (getting patient ready to use the DCA)	0.220
	Types of platforms and devices that can be used	0.212
	Ease of report interpretation	0.197
	Duration of DCA screening (how long it takes)	0.136
Patient and medical assistant	Ease of use for patients	0.330
	Expected patient willingness	0.330
	Ability to integrate into clinic workflow for staff	0.330

ADRD = Alzheimer disease and related dementias; DCA = digital cognitive assessment.

<sup>a</sup> Possible values range from 0 to 1.0, with values within each group summing to 1.0. Higher values indicate greater perceived importance of the criterion relative to others in the group.

each clinic's unique context, to ensure the processes were practical, efficient, and aligned with clinic operations. Weekly reviews and feedback loops during these sprints allowed for continuous refinement and ensured adherence to the Minimum Viable Solution prototype, thereby enhancing fidelity and scalability of the intervention across diverse primary care settings.

## 6. Monitoring Performance

Senior academic and health system leaders, representing primary care, neurology, and geriatrics, met monthly throughout the study period to address high-level gaps or barriers. Agile implementation coaches held a weekly huddle with the physician and staff champion at each clinic to respond quickly to challenges and opportunities. We tracked outcomes through run charts and weekly cross-clinic huddles to monitor patient eligibility and screening completion rates. Cumulative clinic and aggregate dashboard reports were used to discuss implementation progress with clinic and cross-clinic leadership.

## 7. Assessing Organizational Impact

We collaborated with clinic and primary care leaders to evaluate the intervention's impact and to identify any unintended consequences. Clinic-specific performance reports on operational metrics and quality programs continued to perform at targeted performance levels.

## 8. Formulating a Standardized Operating Procedure

We developed a blueprint for implementing DCA screening across different settings based on achieved objectives.

## Workflow Development

During preliminary discussions with primary care leaders about initiating DCA screening, it became clear that managing patients with borderline or positive screenings required a solution that did not increase clinicians' workload. This led to conceptualization of the BHN role. The BHN, a registered nurse, conducted additional assessments, educated patients, provided community resources, assisted primary care clinicians with workups for reversible causes, and facilitated referrals. Clinicians and administrators provided input on BHN scheduling, location, case scope, and communication preferences. The additional assessments conducted included the Montreal Cognitive Assessment, the Patient Health Questionnaire, the Generalized Anxiety Disorder assessment, a review of laboratory and imaging results, and calculation of the anticholinergic burden score and the atherosclerotic cardiovascular disease score.

## Data Collection and Measures

Baseline data were collected for the 12 months before the intervention, and outcomes were monitored during the 12 months after it was implemented. We gathered data from electronic health records, DCA screenings, and weekly cross-clinic huddles. Demographic data were captured to describe

patient characteristics. Screening data included patient eligibility, screening completion, referral completion, and BHN follow-up actions. Huddle and meeting attendance were tracked for clinicians and staff. Electronic health record and screening-related data were entered into a REDCap database (Vanderbilt University and REDCap Consortium) for analysis.

We used the RE-AIM (reach, effectiveness, adoption, implementation, and maintenance) framework to evaluate intervention outcomes across these 5 dimensions.<sup>44,45</sup> This framework accounts for multilevel factors that influence implementation, and provides insights into the generalizability of the intervention, the conditions under which it was implemented, and the outcomes achieved.<sup>44</sup>

## Analysis

We summarized descriptive statistics for baseline clinic characteristics as proportions. Screening completion rates were calculated for each week and the entire 12-month study period.

For the RE-AIM *reach* component, we assessed both clinician reach (defined as the percentage of clinicians engaged in the intervention) and patient reach (defined as the percentage of eligible visits in which DCA screening was completed). For the *effectiveness* component, we assessed selection of the DCA tool, development of a screening workflow, and number of positive screenings resulting in referrals to the BHN. For the *adoption* component, we determined whether clinics achieved full adoption (defined as a cumulative DCA screening completion rate of  $\geq 20\%$  during the 12-month period) or partial adoption (defined as a rate of  $< 20\%$ ). For the *implementation* component, we assessed implementation fidelity, defined as adherence to the Minimum Viable Solution prototype workflow at the end of the study period. For the *maintenance* component, we ascertained continued DCA use 6 and 12 months after implementation, measured by the number of clinics still performing screening and the percentage of referrals to the BHN.

We conducted comparative analysis to explore differences in DCA screening completion rates based on clinician-specific factors, such as hours of sprint sessions attended and clinician engagement during implementation sprints. All statistical analyses were performed using SAS version 9.4 (SAS Institute Inc). Statistical significance was set at  $P < .05$ .

## RESULTS

### Reach

The DCA screening intervention engaged 69 (63.8%) of 108 clinicians across the 7 primary care clinics. Screening was completed in 1,808 (10.8%) of 16,708 eligible visits.

Demographics of the 1,722 unique patients who completed a DCA screening are detailed in an accompanying article.<sup>46</sup> In brief, the patients had a mean (SD) age of 73.5 (6.0) years, 58% were female, and 79% were White. Their median Area Deprivation Index score was 51.

## Effectiveness

After evaluating the 3 DCA tools, stakeholders selected the Linus Health Core Cognitive Evaluation tool,<sup>47</sup> administered and scored using a tablet device and modeled after the paper-and-pencil version of the Mini-Cog tool.<sup>48,49</sup> Although the developed workflows varied by clinic, the Minimum Viable Solution prototype was maintained across clinics, allowing integration of the DCA tool and the BHN into varied clinical settings. Agile implementation permitted clinics to tailor their workflows to either the complete Linus Health Core Cognitive Evaluation or the shorter Linus Health Digital Clock and Recall. The BHN received a total of 447 referrals for further assessment of a positive or borderline screening result. Detailed screening and follow-up results are reported in the accompanying article.<sup>46</sup>

## Adoption

The DCA screening completion rate for unique eligible in-scope patients varied across clinics (Figure 1). Four clinics fully adopted the intervention, achieving a cumulative


DCA screening completion rate of at least 20%: clinic 1 (26.5%), clinic 4 (29.3%), clinic 5 (47.6%), and clinic 6 (31.7%). The other 3 clinics showed partial adoption: clinic 2 (4.1%) and clinic 7 (1.4%), which participated from the start, and clinic 3 (12.0%), which joined 8 months into the intervention.

Fully 87% of early engagers reported readiness to conduct screenings for eligible patients, and early engagers had the highest DCA screening completion rates. There was a moderate, though not significant, correlation of completion rate with the average hours of sprint sessions attended per clinician ( $r = 0.54$ ,  $P = .22$ ) and with clinicians' engagement during implementation sprints ( $r = 0.61$ ,  $P = .15$ ).

## Implementation

All clinics demonstrated fidelity to the Minimum Viable Solution prototype workflow, with consistency of implementation across clinics evident in the run charts. Feedback received during weekly huddles led to iterative adjustments. Table 3 summarizes clinic-specific adaptations.

**Figure 1. Cumulative Percentage of Unique Eligible Patients Within Scope Who Completed a DCA Screening Between June 1, 2022 and May 31, 2023, Overall and by Clinic**



DCA = digital cognitive assessment.

## Maintenance

Twelve months after the intervention began, 5 clinics showed sustained routine use of the DCA tool and the 2 other clinics showed periodic use. Clinicians valued the BHN's support in managing the diagnostic pathway, with ongoing referrals to the BHN still occurring at 12 months. Availability of the BHN was noted as a contributor to maintaining use of the DCA tool in routine practice.

## DISCUSSION

This study evaluated implementation of a DCA tool across diverse primary care clinics using agile implementation. The agile implementation approach enabled successful integration of the DCA tool into routine clinical workflows through structured, iterative, and customizable processes, addressing specific clinic needs.

### Key Findings

#### Workflow Adaptability

Adaptability of the screening workflows to each clinic's unique environment was crucial. This flexibility allowed clinics to overcome initial barriers such as clinicians' lack of readiness and technical difficulties.

#### Clinician Engagement

Targeted engagement sprints proved effective. By identifying clinicians having personas more receptive to agile implementation (early engagers), we tailored strategies to address specific concerns and motivations, leading to improved local adoption across clinics. Future studies should evaluate additional factors related to clinician engagement including revenue, assessing the impact of increased visit complexity or reimbursement of specific Current Procedural Terminology billing codes in primary care.<sup>50</sup>

#### BHN Role

The BHN played a key role in managing positive and borderline screenings and reducing related primary care clinician workload. By facilitating additional assessments and ensuring appropriate follow-up, the BHN supported patients and caregivers in the diagnostic pathway and added considerable value

for clinicians. Future research should evaluate this role's effectiveness and reimbursement mechanisms for sustainability.

#### Patient Reach and Screening Completion

The intervention reached a substantial number of patients, with 1,808 DCA screenings completed during 12 months. Despite an overall screening completion rate of only 10.8%, the progressive increase in rate throughout the period demonstrated growing and sustained integration of the DCA tool into routine clinic operations at several clinics.

#### Cost-Effectiveness

Beyond the incremental costs of adding DCA screening to our existing primary care workforce, implementation required incremental resources including the BHN, agile implementation coaching and facilitation, and support for data collection and analysis. As part of the Davos Flagship program, DCA expenses were provided in-kind. The growing prevalence of dementia and its associated economic burden<sup>51,52</sup> underscores the need to evaluate the cost-effectiveness of screening interventions. Previous studies suggest that dementia screening programs can be cost-effective.<sup>4-6</sup> Future research should assess economic sustainability while delivering measurable clinical benefits.

### Challenges

#### Initial Technical Barriers

Initial technical challenges, including DCA tool integration, were resolved through iterative adjustments and feedback mechanisms.

#### Staffing Shortages

Staffing shortages, particularly vacant medical assistant positions during the study period, impeded initial implementation efforts. Addressing these shortages through strategic workforce planning is essential for sustaining screening workflows and should be considered in future studies. Although the BHN supported patient assessment after a DCA screening, medical assistants and clinicians still faced added burden from additional screening tasks. Strategic evaluation of primary care workforce ratios and reimbursement models at the health system, state, and national levels is necessary to address the broader impact of primary care-based screenings.

#### Influence of Recommendations

The lack of a US Preventive Services Task Force endorsement for routine cognitive screening in older adults<sup>53</sup> presented a challenge to clinician engagement. Despite recommendations from other professional organizations, such as the American Academy of Neurology, which advocates for annual cognitive screenings,<sup>54</sup> clinician ambivalence persisted. In addition, patients' reluctance to report

**Table 3. Clinic-Specific Workflow Adaptations**

Development and dissemination of succinct training videos for the independent instruction of medical assistants in the administration of DCA
Extension of the BHN's duties to include the calculation of anticholinergic burden scores
Establishment of a backup system to ensure BHN's reliable communications of results
Customization of screening workflows, using the digital tool's configurability, offering options from a comprehensive 15-minute DCA inclusive of a lifestyle survey to a brief 3-minute assessment
Enhancements to the result display, to improve interpretation of results for clinicians
Streamlining of the referral process to geriatrics, neurology, and research studies

BHN = brain health navigator; DCA = digital cognitive assessment.

memory concerns<sup>54</sup> highlights the need for more robust evidence on the benefits of early cognitive assessment. A key component of guideline-supported screening is its practical implementation in routine care. This study offers insights into workflow integration and adoption of DCA tools through successful implementation.

### Implications for Practice

Adoption of an agile implementation framework for implementing interventions in health care settings has demonstrated considerable potential to enhance both effectiveness and sustainability.<sup>24,25,28,31-34,55</sup> The agile implementation model highlights the critical importance of adaptability and stakeholder engagement in successful implementation.

### Limitations

This study has several limitations. First, the nonrandom selection of clinics and voluntary participation of clinicians may have introduced selection bias, affecting the generalizability of the results. Our study used quality improvement methods, which place greater emphasis on ongoing measurement in a real-world context but entail less emphasis on bias, less control of confounding variables, and less stringent population parameters than traditional research, consistent with other quality improvement methods.<sup>36</sup> Second, variations in practice type, staffing, and available physical space for activities could have influenced implementation outcomes. Third, the 12-month study period limits evaluation of the intervention's long-term sustainability and impact. Future studies should explore randomized designs and conduct cost-effectiveness analyses to ensure scalability and sustainability.

## CONCLUSIONS

This study demonstrates successful implementation of a DCA tool in primary care using agile implementation principles. Key factors for adoption and sustainability included tailored DCA screening workflows and the integration of a BHN role, which contributed to a screening completion rate of 10.8%. Addressing staffing shortages and establishing sustainable revenue models are critical for ongoing implementation.

Our findings underscore the value of iterative improvement and stakeholder engagement in implementing health interventions. Future research should extend follow-up periods, expand to a broader range of clinical settings, and evaluate cost-effectiveness and the long-term impact on patient care and the health care system.

 [Read or post commentaries in response to this article.](#)

**Key words:** dementia screening; primary health care; digital cognitive assessment; quality improvement; workflow; implementation science; health informatics; cognitive dysfunction; Alzheimer's disease; neuropsychologic tests; mild cognitive impairment; practice-based research; clinical practice patterns; organizational innovation

Submitted June 20, 2024; submitted, revised, January 14, 2025; accepted February 4, 2025.

**Funding support:** This study was funded by the Davos Alzheimer's Collaborative Healthcare System Preparedness Program.

**Disclaimer:** Linus Health did not have any role in the design of the study; the conduct of the study; the collection, management, analysis, or interpretation of the data; the preparation of the manuscript; or the decision to submit the manuscript for publication.

**Acknowledgments:** This Early Detection program was made possible by the support of the Davos Alzheimer's Collaborative Healthcare System Preparedness program. Linus Health provided in-kind use of the Linus Health Core Cognitive Evaluation and Digital Clock Recall, and technical support for this project.

## References

- Alzheimer's disease facts and figures. *Alzheimers Dement*. 2024;20(5):3708-3821. doi:[10.1002/alz.13809](https://doi.org/10.1002/alz.13809)
- Cai C, Gaffney A, McGregor A, et al. Racial and ethnic disparities in outpatient visit rates across 29 specialties. *JAMA Intern Med*. 2021;181(11):1525-1527. doi:[10.1001/jamainternmed.2021.3771](https://doi.org/10.1001/jamainternmed.2021.3771)
- Chin AL, Negash S, Hamilton R. Diversity and disparity in dementia: the impact of ethnoracial differences in Alzheimer disease. *Alzheimer Dis Assoc Disord*. 2011;25(3):187-195. doi:[10.1097/WAD.0b013e318211c6c9](https://doi.org/10.1097/WAD.0b013e318211c6c9)
- Saito E, Nakamoto BK, Mendez MF, Mehta B, McMurtray A. Cost effective community based dementia screening: a Markov model simulation. *Int J Alzheimers Dis*. 2014;2014:103138. doi:[10.1155/2014/103138](https://doi.org/10.1155/2014/103138)
- Yu SY, Lee TJ, Jang SH, Han JW, Kim TH, Kim KW. Cost-effectiveness of nationwide opportunistic screening program for dementia in South Korea. *J Alzheimers Dis*. 2015;44(1):195-204. doi:[10.3233/jad-141632](https://doi.org/10.3233/jad-141632)
- Zhang Y, Kivipelto M, Solomon A, Wimo A. Cost-effectiveness of a health intervention program with risk reductions for getting demented: results of a Markov model in a Swedish/Finnish setting. *J Alzheimers Dis*. 2011;26(4):735-744. doi:[10.3233/JAD-2011-110065](https://doi.org/10.3233/JAD-2011-110065)
- Akushevich I, Kravchenko J, Yashkin A, Doraiswamy PM, Hill CV; Alzheimer's Disease and Related Dementia Health Disparities Collaborative Group. Expanding the scope of health disparities research in Alzheimer's disease and related dementias. *Alzheimers Dement (Amst)*. 2023;15(1):e12415. doi:[10.1002/dad2.12415](https://doi.org/10.1002/dad2.12415)
- Budd Haeberlein S, Aisen PS, Barkhof F, et al. Two randomized phase 3 studies of aducanumab in early Alzheimer's disease. *J Prev Alzheimers Dis*. 2022;9(2):197-210. doi:[10.14283/jpad.2022.30](https://doi.org/10.14283/jpad.2022.30)
- Sims JR, Zimmer JA, Evans CD, et al; TRAILBLAZER-ALZ 2 Investigators. Donanemab in early symptomatic Alzheimer disease: The TRAILBLAZER-ALZ 2 Randomized Clinical Trial. *JAMA*. 2023;330(6):512-527. doi:[10.1001/jama.2023.13239](https://doi.org/10.1001/jama.2023.13239)
- Swanson CJ, Zhang Y, Dhadda S, et al. A randomized, double-blind, phase 2b proof-of-concept clinical trial in early Alzheimer's disease with lecanemab, an anti-A protofibril antibody. *Alzheimers Res Ther*. 2021;13(1):80. doi:[10.1186/s13195-021-00813-8](https://doi.org/10.1186/s13195-021-00813-8)
- van Dyck CH, Swanson CJ, Aisen P, et al. Lecanemab in early Alzheimer's disease. *N Engl J Med*. 2023;388(1):9-21. doi:[10.1056/NEJMoa2212948](https://doi.org/10.1056/NEJMoa2212948)
- de Levante Raphael D. The knowledge and attitudes of primary care and the barriers to early detection and diagnosis of Alzheimer's disease. *Medicina (Kaunas)*. 2022;58(7):906. doi:[10.3390/medicina58070906](https://doi.org/10.3390/medicina58070906)
- Amjad H, Roth DL, Sheehan OC, Lyketsos CG, Wolff JL, Samus QM. Underdiagnosis of dementia: an observational study of patterns in diagnosis and awareness in US older adults. *J Gen Intern Med*. 2018;33(7):1131-1138. doi:[10.1007/s11606-018-4377-y](https://doi.org/10.1007/s11606-018-4377-y)
- Bernstein A, Rogers KM, Possin KL, et al. Dementia assessment and management in primary care settings: a survey of current provider practices in the United States. *BMC Health Serv Res*. 2019;19(1):919. doi:[10.1186/s12913-019-4603-2](https://doi.org/10.1186/s12913-019-4603-2)
- Lang L, Clifford A, Wei L, et al. Prevalence and determinants of undetected dementia in the community: a systematic literature review and a meta-analysis. *BMJ Open*. 2017;7(2):e011146. doi:[10.1136/bmjopen-2016-011146](https://doi.org/10.1136/bmjopen-2016-011146)

16. Liu Y, Jun H, Becker A, Wallick C, Mattke S. Detection rates of mild cognitive impairment in primary care for the United States Medicare population. *J Prev Alzheimers Dis*. 2024;11(1):7-12. doi:[10.14283/jpad.2023.131](https://doi.org/10.14283/jpad.2023.131)
17. Borson S, Small GW, O'Brien Q, Morrello A, Boustani M. Understanding barriers to and facilitators of clinician-patient conversations about brain health and cognitive concerns in primary care: a systematic review and practical considerations for the clinician. *BMC Prim Care*. 2023;24(1):233. doi:[10.1186/s12875-023-02185-4](https://doi.org/10.1186/s12875-023-02185-4)
18. Dumas A, Destrebecq F, Esposito G, Suchonova D, Steen Frederiksen K. Rethinking the detection and diagnosis of Alzheimer's disease: outcomes of a European Brain Council project. *Aging Brain*. 2023;4:100093. doi:[10.1016/j.nbas.2023.100093](https://doi.org/10.1016/j.nbas.2023.100093)
19. Koch T, Illiffe S; EVIDEM-ED project. Rapid appraisal of barriers to the diagnosis and management of patients with dementia in primary care: a systematic review. *BMC Fam Pract*. 2010;11(1):52. doi:[10.1186/1471-2296-11-52](https://doi.org/10.1186/1471-2296-11-52)
20. Mansfield E, Noble N, Sanson-Fisher R, Mazza D, Bryant J. Primary care physicians' perceived barriers to optimal dementia care: a systematic review. *Gerontologist*. 2019;59(6):e697-e708. doi:[10.1093/geront/gny067](https://doi.org/10.1093/geront/gny067)
21. Podhorna J, Winter N, Zobelein H, Perkins T. Alzheimer's treatment: real-world physician behavior across countries. *Adv Ther*. 2020;37(2):894-905. doi:[10.1007/s12325-019-01213-z](https://doi.org/10.1007/s12325-019-01213-z)
22. Dham P, McAiney C, Saperson K, et al. Impact of integrated care pathways within the framework of collaborative care on older adults with anxiety, depression, or mild cognitive impairment. *Am J Geriatr Psychiatry*. 2022;30(7):834-847. doi:[10.1016/j.jagp.2022.01.010](https://doi.org/10.1016/j.jagp.2022.01.010)
23. Olswang LB, Prelock PA. Bridging the gap between research and practice: implementation science. *J Speech Lang Hear Res*. 2015;58(6):S1818-S1826. doi:[10.1044/2015\\_JSLHR-L-14-0305](https://doi.org/10.1044/2015_JSLHR-L-14-0305)
24. Boustani M, Alder CA, Solid CA. Agile implementation: a blueprint for implementing evidence-based healthcare solutions. *J Am Geriatr Soc*. 2018;66(7):1372-1376. doi:[10.1111/jgs.15283](https://doi.org/10.1111/jgs.15283)
25. Boustani M, Azar J, Solid C. *Agile Implementation: A Model for Implementing Evidence-Based Healthcare Solutions Into Real-World Practice to Achieve Sustainable Change*. Morgan James Publishing; 2020.
26. Chamorro-Koc M. Prototyping for healthcare innovation. In: Miller E, Winter A, Chari S, eds. *How Designers Are Transforming Healthcare*. Springer Nature Singapore; 2024:103-117.
27. Caballero L, Moreno AM, Seffah A. Persona as a tool to involving human in Agile methods: contributions from HCI and marketing. In: *Human-Centered Software Engineering*. Proceedings of the 5th IFIP WG 13.2 International Conference, HCSE 2014, Paderborn, Germany, September 16-18, 2014. Springer Berlin Heidelberg; 2014:283-290.
28. Boustani MA, Van der Marck MA, Adams N, et al. Developing the agile implementation playbook for integrating evidence-based health care services into clinical practice. *Acad Med*. 2019;94(4):556-561. doi:[10.1097/ACM.0000000000002497](https://doi.org/10.1097/ACM.0000000000002497)
29. Puspitasari I, Nuzulita N, Hsiao C-S. Agile user-centered design framework to support the development of e-health for patient education. In: Lee R, ed. *Computer and Information Science and Engineering: Volume 16*. Springer Nature Switzerland; 2024:131-144.
30. Dobre J, Harrington C, Herout J, et al. Rapid heuristic evaluation: ensuring fast and reliable usability support. *Proc Hum Factors Ergon Soc Annu Meet*. 2017;61(1):610-614. doi:[10.1177/1541931213601638](https://doi.org/10.1177/1541931213601638)
31. Kitzmiller R, Hunt E, Sproat SB. Adopting best practices: "Agility" moves from software development to healthcare project management. *Comput Inform Nurs*. 2006;24(2):75-82. doi:[10.1097/00024665-200603000-00005](https://doi.org/10.1097/00024665-200603000-00005)
32. Desai M, Tardif-Douglin M, Miller I, et al. Implementation of Agile in healthcare: methodology for a multisite home hospital accelerator. *BMJ Open Qual*. 2024;13(2):e002764. doi:[10.1136/bmjog-2024-002764](https://doi.org/10.1136/bmjog-2024-002764)
33. Saleh HH, Abbas ZA, Latif N, Khalil ZT. Agile management in healthcare improving patient outcomes through flexibility and responsiveness. *J Ecohumanism*. 2024;3(5):633-649. doi:[10.62754/joe.v3i5.3928](https://doi.org/10.62754/joe.v3i5.3928)
34. Sindhvani R, Singh PL, Prajapati DK, Iqbal A, Phanden RK, Malhotra V. Agile system in health care: literature review. In: Shanker K, Shankar R, Sindhvani R, eds. *Advances in Industrial and Production Engineering. Lecture Notes in Mechanical Engineering*. Springer; 2019. doi:[10.1007/978-981-13-6412-9\\_61](https://doi.org/10.1007/978-981-13-6412-9_61)
35. Carter N, Valaitis RK, Lam A, Feather J, Nicholl J, Cleghorn L. Navigation delivery models and roles of navigators in primary care: a scoping literature review. *BMC Health Serv Res*. 2018;18(1):96. doi:[10.1186/s12913-018-2889-0](https://doi.org/10.1186/s12913-018-2889-0)
36. Perla RJ, Provost LP, Murray SK. Sampling considerations for health care improvement. *Qual Manag Health Care*. 2014;23(4):268-279. doi:[10.1097/qmh.0000000000000042](https://doi.org/10.1097/qmh.0000000000000042)
37. Finkelstein JA, Brickman AL, Capron A, et al. Oversight on the borderline: quality improvement and pragmatic research. *Clin Trials*. 2015;12(5):457-466. doi:[10.1177/17407745155597682](https://doi.org/10.1177/17407745155597682)
38. Pannick S, Sevdalis N, Athanasiou T. Beyond clinical engagement: a pragmatic model for quality improvement interventions, aligning clinical and managerial priorities. *BMJ Qual Saf*. 2016;25(9):716-725. doi:[10.1136/bmjqs-2015-004453](https://doi.org/10.1136/bmjqs-2015-004453)
39. Kind AJH, Buckingham WR. Making neighborhood-disadvantage metrics accessible - The Neighborhood Atlas. *N Engl J Med*. 2018;378(26):2456-2458. doi:[10.1056/NEJMp1802313](https://doi.org/10.1056/NEJMp1802313)
40. University of Wisconsin School of Medicine and Public Health. Neighborhood Atlas. Area deprivation index. Accessed Mar 2024. <https://www.neighborhoodatlas.medicine.wisc.edu/>
41. Nasa P, Jain R, Juneja D. Delphi methodology in healthcare research: how to decide its appropriateness. *World J Methodol*. 2021;11(4):116-129. doi:[10.5662/wjm.v11.i4.116](https://doi.org/10.5662/wjm.v11.i4.116)
42. Davos Alzheimer's Collaborative. The Davos Alzheimer's Collaborative Launches Innovative Early Detection Effort: Seven Flagship Pilot Sites Aim to Measurably Increase Timely, Accurate Diagnosis of Alzheimer's. Published May 3, 2022. Accessed Mar 31, 2025. <https://www.davosalzheimerscollaborative.org/news-press/https://www.davosalzheimerscollaborative.org/news-and-press/davos-alzheimers-collaborative-launches-innovative-early-detection-effort>
43. Mazak A, Kratzwald B. The Weighted Decision Matrix: tracking design decisions in service compositions. In: *2014 IEEE 7th International Conference on Service-Oriented Computing and Applications*. Matsue, Japan; 2014:169-174. doi:[10.1109/SOCA.2014.55](https://doi.org/10.1109/SOCA.2014.55)
44. Glasgow RE, Harden SM, Gaglio B, et al. RE-AIM planning and evaluation framework: adapting to new science and practice with a 20-year review. *Front Public Health*. 2019;7:64. doi:[10.3389/fpubh.2019.00064](https://doi.org/10.3389/fpubh.2019.00064)
45. Holtrop JS, Estabrooks PA, Gaglio B, et al. Understanding and applying the RE-AIM framework: clarifications and resources. *J Clin Transl Sci*. 2021;5(1):e126. doi:[10.1017/cts.2021.789](https://doi.org/10.1017/cts.2021.789)
46. Fowler NR, Hammers DB, Perkins AJ, et al. Feasibility and acceptability of implementing a digital cognitive assessment for Alzheimer disease and related dementias in primary care. *Ann Fam Med*. 2025;23(3):online. doi:[10.1370/afm.240293](https://doi.org/10.1370/afm.240293)
47. Libon DJ, Matusz EF, Cosentino S, et al. Using digital assessment technology to detect neuropsychological problems in primary care settings. *Front Psychol*. 2023;14:1280593. doi:[10.3389/fpsyg.2023.1280593](https://doi.org/10.3389/fpsyg.2023.1280593)
48. Seitz DP, Chan CC, Newton HT, et al. Mini-Cog for the diagnosis of Alzheimer's disease dementia and other dementias within a primary care setting. *Cochrane Database Syst Rev*. 2018;2(2):CD011415. doi:[10.1002/14651858.CD011415.pub2](https://doi.org/10.1002/14651858.CD011415.pub2)
49. Seitz DP, Chan CC, Newton HT, et al. Mini-Cog for the detection of dementia within a primary care setting. *Cochrane Database Syst Rev*. 2021;7(7):CD011415. doi:[10.1002/14651858.CD011415.pub3](https://doi.org/10.1002/14651858.CD011415.pub3)
50. Alzheimer's Association. Cognitive Assessment and Care Planning Services: Alzheimer's Association Expert Task Force Recommendations and Tools for Implementation. Care planning visit. Accessed Feb 12, 2025. <https://www.alz.org/careplanning/downloads/cms-consensus.pdf>
51. Cantarero-Prieto D, Leon PL, Blazquez-Fernandez C, Juan PS, Cobo CS. The economic cost of dementia: a systematic review. *Dementia (London)*. 2020;19(8):2637-2657. doi:[10.1177/1471301219837776](https://doi.org/10.1177/1471301219837776)
52. Wimo A, Seeher K, Cataldi R, et al. The worldwide costs of dementia in 2019. *Alzheimers Dement*. 2023;19(7):2865-2873. doi:[10.1002/alz.12901](https://doi.org/10.1002/alz.12901)
53. Owens DK, Davidson KW, Krist AH, et al.; US Preventive Services Task Force. Screening for Cognitive Impairment in Older Adults: US Preventive Services Task Force Recommendation Statement. *JAMA*. 2020;323(8):757-763. doi:[10.1001/jama.2020.0435](https://doi.org/10.1001/jama.2020.0435)
54. Foster NL, Bondi MW, Das R, et al. Quality improvement in neurology: mild cognitive impairment quality measurement set. *Neurology*. 2019;93(16):705-713. doi:[10.1212/WNL.0000000000008259](https://doi.org/10.1212/WNL.0000000000008259)
55. Summanwar D, Ropert C, Barton J, et al. Agile implementation of alcohol screening in primary care. *BMC Prim Care*. 2024;25(1):251. doi:[10.1186/s12875-024-02500-7](https://doi.org/10.1186/s12875-024-02500-7)