

Online Supplementary Material

Gaynes BN, Rush A, Trivedi M, et al. Major depression symptoms in primary care and psychiatric care settings: a cross-sectional analysis. *Ann Fam Med.* 2007;5(2):126-134.

<http://www.annfammed.org/cgi/content/full/5/2/126/DC1>

Supplemental Table. Percentage of Patients With Specific Depressive Symptoms (IDS-C30 by ROA)				
Selected IDS-C30 (ROA) Items	Recruitment Setting		Unadjusted P Value	Adjusted P Value*
	Primary Care n = 1,063 (41.8%)	Specialty Care n = 1,478 (58.2%)		
Essential features				
Depressed mood (sad)	96.6	97.9	.0486	.0531
Anhedonia: involvement (interest in activities)	80.6	85.7	.0015	.0025
Core features				
Appetite/weight change				
Appetite decreased	42.7	44.3	.4524	.4992
Appetite increased	23.3	19.8	.0475	.2556
Weight decrease	29.2	29.3	.9740	.9245
Weight increase	25.0	22.3	.1278	.6157
Sleep disturbance				
Sleep onset insomnia	67.1	66.8	.9029	.5168
Mid-nocturnal insomnia	83.0	77.2	.0008	.5547
Early morning insomnia	52.4	50.3	.3234	.0806
Hypersomnia	21.3	26.6	.0045	.7476
Psychomotor change				
Slowing	62.1	61.8	.8845	.4531
Agitation	60.7	64.5	.0682	.0188
Fatigue/loss of energy (energy/fatigability)				
Fatigue/loss of energy (energy/fatigability)	89.9	88.4	.2723	.9027
Worthless/guilt (outlook-self)				
Worthless/guilt (outlook-self)	76.6	82.5	.0006	.0755
Decreased concentration (concentration/decision making)				
Decreased concentration (concentration/decision making)	88.5	91.8	.0087	.0084
Suicidal ideation				
Suicidal ideation	42.8	51.4	<.0001	<.0001

Note: Percentage for which the symptom was rated as present (> 1 on the item).
 IDS-C30 = 30-item Inventory of Depressive Symptomatology, Clinician Rating; ROA = research outcome assessor.
 *Adjusted for age, sex, education, employment status, marital status, ethnicity, length of current episode, and Cumulative Illness Rating Scale (total score).