

Supplemental materials for:

Shires DA, Stroumsa D, Jaffee KD, Woodford MR. Primary care clinicians' willingness to care for transgender patients. *Ann Fam Med*. 2018;16(6):555-558.

Supplemental Appendix 1. Multivariable Logistic Regression: Factors Predicting Primary Care Clinicians' Willingness to *Provide Routine Care* to Transgender Patients (N=140)^a

Variable ^b	AOR	95% CI	p
Age	0.96	0.91-0.99	.039
Continent of origin: North America/Caribbean	2.52	0.76-8.30	.129
Family medicine (vs. Internal medicine)	1.32	0.39-4.43	.656
Ever met transgender person	1.60	0.40-6.34	.505
Cared for transgender patient in past 5 years	2.25	0.57-8.88	.247
Empathy	1.31	0.91-1.86	.143
Transphobia	0.82	0.48-1.42	.479
Lack of training on transgender health	0.53	0.52-5.38	.593
Lack of exposure to transgender patients	0.63	0.11-3.64	.606
Lack of knowledge among staff	0.41	0.08-2.19	.299
Lack of familiarity with guidelines	4.48	0.51-39.22	.175

^a $\chi^2(11, N=140) = 26.19, p=.006$, Nagelkerke $R^2 = .306$

^b *Feels capable of providing routine care* was not included in the model due to a lack of variability and resulting instability

Supplemental Appendix 2. Multivariable Logistic Regression: Factors Predicting Primary Care Clinicians' Willingness to *Provide Pap Tests for Transgender Men* (N=140)^a

Variable	AOR	95% CI	p
Age	0.97	0.94-1.01	.123
Continent of origin: North America/Caribbean	1.04	0.35-3.09	.945
Family medicine (vs. Internal medicine)	5.08	1.35-19.16	.016
Ever met transgender person	4.11	1.13-14.96	.032
Cared for transgender patient in past 5 years	1.14	0.33-3.94	.832
Empathy	1.18	0.84-1.67	.339
Transphobia	0.54	0.32-0.91	.020
Lack of training on transgender health	1.05	0.16-3.78	.964
Lack of exposure to transgender patients	0.77	0.16-3.78	.746
Lack of knowledge among staff	0.88	0.19-4.09	.872
Lack of familiarity with guidelines	0.59	0.09-3.87	.585
Feels capable of providing routine care	2.11	0.72-6.19	.172

^a $\chi^2(12, N=140) = 38.83, p<.001$, Nagelkerke $R^2 = .376$