

Supplemental materials for:

Patchett-Marble R. A Staff- and protocol-driven system to apply urine drug testing in primary care. *Ann Fam Med*. 2020;18(6):563.

Submitted April 26, 2019; submitted, revised, March 16, 2020; accepted March 24, 2020.

Funding support: Supported by the Northern Ontario Academic Medicine Association (NOAMA) AHSC AFP Innovation Fund award.

Previous presentations: Family Medicine Forum; November 2018; Toronto, Ontario, Canada; and Association of Family Health Teams of Ontario Annual Conference; September, 2019; Toronto, Ontario, Canada.

References

1. Gomes T, Pasricha S, Martins D, Greaves S. Behind the Prescriptions: A Snapshot of Opioid Use Across All Ontarians. Toronto, Canada: Ontario Drug Policy Research Network; 2017. Submitted April 26, 2019; submitted, revised, March 16, 2020; accepted March 24, 2020.

2. Argoff CE, Alford DP, Fudin J, et al. Rational urine drug monitoring in patients receiving opioids for chronic pain: consensus recommendations. *Pain Med*. 2018;19(1):97-117. 10.1093/pm/pnx285.

Supplemental Figures 1-3:

IF RISKS > BENEFITS
•SUPPORT THE PATIENT
•TAPER AND DISCONTINUE OPIOIDS

**OPIOID ADDICTION:
OPIOID AGONIST TREATMENT**

**DAILY DISPENSING
2-4 URINE TESTS PER MONTH**

**SHORT DISPENSING INTERVALS,
6-7 URINE TESTS PER YEAR**

**SHORT DISPENSING INTERVALS,
2-3 URINE TESTS PER YEAR**

**SHORT DISPENSING INTERVALS,
1-2 URINE TESTS PER YEAR**

**LONG DISPENSING INTERVALS,
1-2 URINE TESTS PER YEAR**

.....▶ = CONCERN IDENTIFIED FOR CLINICAL INSTABILITY ▶ = DEMONSTRATED CLINICAL STABILITY





